

## **REVIEW OF USE OF ALTERNATIVES TO METHYL BROMIDE FOR HORTICULTURAL CROPS IN ASIA**

**J.G.M. Vos\* and G.S. Lim**

Methyl bromide is a broad spectrum pesticide, principally used as a fumigant. On soil, it is used as a pre-plant soil fumigant in locations where a broad complex of soil-borne pests limit the production of certain crops, particularly when they are repeatedly grown on the same land. In SE Asia and the Pacific, the use of methyl bromide as a soil sterilant is only about 6 % of all methyl bromide used (UNDP, 1995). Alternative methods to control soil pests are widespread. Aware of the negative side effects of pesticides, alternative soil pest control practices with reduced chemical inputs are being suggested. Soil pest control practices include biological control by means of natural enemies and integrated methods, such as integrated pest management (IPM) or integrated crop management (ICM). This paper documents a number of cases of alternative soil pest control methods with reduced chemical inputs in commercial use for horticultural crops in Asia. In addition, alternatives that have been found effective in field and/or greenhouse trials are reviewed.

Cases of alternative soil pest control methods used commercially for horticultural crops on modern as well as traditional farms in Asia are listed in Table 1. The methods range from relatively new techniques, such as substrates and solarisation, to improved methods of traditional techniques, such as composting. There is now greater appreciation and respect for low-input, traditional agricultural systems, and the methods that are employed by traditional farmers are recognised as often being more efficient and logical than previously thought. In fact, to manage nematodes, integrated systems are being developed that incorporate or consider many alternative small-scale farmer practices (Bridge, 1996). Most of the identified techniques can be applied in intensive horticultural production.

In addition to the cases of commercially applied alternative soil pest control methods described in the previous section, there are numerous reports of IPM experimental efforts and successes in many parts of the world, and farmers would benefit from further verification on a larger scale at the farm level. Table 2 lists cases of experimental use of soil pest control methods that could be used as elements of an IPM / ICM approach to successfully replace methyl bromide, with a general indication of applicability.

This review shows that alternative methods to manage soil-borne pests are widespread on various crops in Asia. The choice of method(s) varies to suit the cropping system and the environment. It is concluded that practices do exist to manage soil-borne pests successfully in ecologically sound ways to achieve sustainable crop production.

It should be stressed that, in addition to the documented cases given here, many other (cultural) practices are widespread throughout the Asian region which are applied for various reasons. These, integrated with other crop management measures contribute to the management of soil pests in general, such as sanitation, various methods of tillage (e.g. repeated summer ploughing), crop rotation, balanced fertilising, regulation of irrigation, optimising planting time, choice of seed / planting material, planting distance and habitat management. IPM integrates the available pest control methods to achieve a farmer's most effective, economical, and sustainable combination for a particular local situation. There are now many IPM success cases reported at the farm level for a variety of crops, including horticultural crops. In Asia, all of these success cases contradict the claim that methyl bromide would be needed to grow commercial horticultural crops successfully and economically.

**Table 1: Cam using alternative practices for the effective control of soil pests commercially.**

<u>crop(s)</u>	<u>Method</u>	<u>Country/ies</u>	<u>Source(s) of information</u>
1 various horticultural crops	Composting and using compost	India Malaysia Philippines South Korea Thailand Elsewhere	IIBC staff Harris (1996) FOAM (1996)
2 Shallot Hot popper Tomato Other vegetables	Flooding of fields	Indonesia Malaysia Vietnam Elsewhere	IIBC staff Vos et al. (1993) Sariah and Tanaka (1994) Vu (1990)
3 Strawberry Tomato Pepper Leafy vegetables	Cultivation on substrates	Malaysia Indonesia Singapore	IIBC staff 'Various private growers
4 Cabbage Solanaceous crops	Application of lime	Malaysia Thailand Philippines Elsewhere	IIBC staff Vattanatengum (1990)
5 Pepper Tomato Watermelon	Application of mulch	Taiwan Indonesia Malaysia Vietnam	IIBC staff Vos and Sumarni (1997) Vu (1990)
6 Cabbage	Treatment of seed	Countries in Asia India Bangladesh	IIBC staff Vattanatangum (1990) IIBC staff Choudhury and Hoque (1982)
7 Various horticultural crops	Burning of top-soil	India	IIBC staff
8 Various horticultural crops	Application of neem cake	India	IIBC staff Alam (1991)
9 Various horticultural crops	Mixed cropping with marigold	India Vietnam	IIBC staff Khan et al. (1971)
10 various horticultural crops	Tillage of soil	Vietnam Bangladesh India	IIBC staff Vu (1990) Hossain (1990) Prasad and Chawla (1991)
11 Cucumber Eggplant Tomato	Grafting on resistant root stocks	China Japan	IIBC staff Qiu (1990)
12 Various horticultural crops	Use of resistant cultivars.	Japan Malaysia Elsewhere	IIBC staff Takeuchi (1990)
13 Various horticultural crops	Solarisation in greenhouses	Japan	Horiuchi (1991)
14 Potato	Increase of hilling and irrigation	Australia	Osmelak (1990)
15 Tomato	Rotation of	Bangladesh	Hossain (1990)
16 Various horticultural crops	Solarisation of seed beds	China	Qiu (1990)
17 Tomato Eggplant Other vegetables	Adjusting of planting time	India	Sethi and Gaur (1986)

Table 2: **Alternative soil pest control methods found effective in tests or trials.**

<u>Crop(s)</u>	<u>Method</u>	<u>Country/ies</u>	<u>Source(s) of information</u>	<u>Applicability</u>
1 Tomato	Resistance against bacterial wilt	Indonesia	Purwati and Hanudin (1995)	Regions with equal wilt strains
2 Cabbage	Solarisation	Elsewhere	Boteng (1990)	Regions with hot sunny climates
Onion		Philippines	Porter and Merriman (1985)	
3 Eggplant	Biological control using <i>Talaromyces flavus</i>	Australia		Fields and greenhouses
4 Cumin	Solarisation, summer irrigation, amendments	USA	Marois et al. (1982)	Hot and regions
Clusterbean		India	Lodha (1995)	
5 Tomato	Integration of biological and chemical control	Israel	Elad et al. (1994)	Fields and greenhouses with <i>Botrytis cinerea</i>
Strawberry				Tomato Wowing areas
6 Tomato	Resistance to rootknot nematode	Argentina	Gallardo (1988)	Greenhouse lettuce crops
7 Lettuce	Biological control using <i>Trichoderma viride</i>	Elsewhere	Bedlan (1988)	Vegetable crops in greenhouses and fields
8 Cucumber	Integration of biological and cultural control	Austria	Kobayashi (1991)	
Tomato		Japan		
9 Tomato	Use of trap crops	India	Rangaswamy and Reddy (1993)	Greenhouses and fields
10 Tomato	Grafting on resistant rootstocks	Switzerland	Dufour and Taillens (1994)	Fields and greenhouses
Cucumber		Japan		
Eggplant		China	Kuwata et al. (1994)	
Pepper		Italy	Lu et al. (1992)	
		Korea	Mona et al. (1992)	
11 Banana	Hot water treatment	?	Choe (1989)	Small-hold farms
			Prasad and Reddy (1994)	
12 Tea	Soil sterilisation for young tea plants	India	Rao (1976)	Small-hold farms

## References:

- Alam, M.M. (1991)  
Control of plant-parasitic nematodes with oilseed cakes on some vegetables in field Pakistani Journal of Nematology 9: p. 21-30
- Bedlan, G. (1988)  
Bekämpfungsmöglichkeiten der Schwarzfaule mit Hilfe des Antagonisten *Trichoderma viride* Pers. an Salat im Freiland. Pflanzenschutz 4: p. 8-10
- Boteng, M.D.C. (1990) Control of clubroot (*Plasmodiophora brassicae*) on cabbage (*Brassica oleracea*). B.S. thesis, Benguet State University, La Trinidad, Benguet, Philippines: 27 pp.
- Bridge, J. (1996) Nematode management in sustainable and subsistence agriculture. Annual Review of Phytopathology 34: p. 201-225.
- Choe, J.S. (1989)  
Phytophthora blight of green pepper in Korea. Extension bulletin ASPAC, Food and fertilizer technology centre 302: p. 18-25
- Choudhury, B.C. and Hoque, M.O. (1982)  
Control of root-knot nematode in brinjal seed beds by heat treatments. Bangladesh Journal of Agricultural Research 7: p. 55-57
- Dufour, L. and Taillens, J. (1994)  
Le greffage: une methode de lutte efficace contre le fusariose vasculaire de la tomate et la pourriture noire des racines du concombre. Revue Suisse de Viticulture 26(2): p. 85-88
- Elad, Y., Shtienberg, D. and Niv, A. (1994) *Trichoderma harzianum* T39 integrated with fungicides: improved biocontrol of grey mould. Brighton crop protection conference -pests and diseases 1994: p. 1109-1114
- Gallardo, G.S. (1988) Breeding of processing tomatoes with genetic resistance to rootknot nematodes (*Meloidogyne Incognita*). Proceedings of the international symposium on integrated management practices in tomato and pepper production in the tropics, Tainan, Taiwan. AVRDC. p. 107-112
- Harris, P.J.C. (1996) An assessment of the potential contribution of organic expertise and techniques to the development of sustainable farming systems for the Eastern India Rainfed Farming Project. Consultancy report, Henry Doubleday Research Association, UK: 45 pp.
- Horiuchi, S. (1991)  
Solarisation for greenhouse crops in Japan. In: DeVay, J.E., Stapleton, J.J. and Elmore, C.L. (eds) Soil solarisation. Proceedings of the first international conference on soil solarisation. FAO plant production and protection paper 109: p. 16-27
- Hossain, Q.T. (1990)  
Status and management of vegetable pests in Bangladesh. In: Lim, G.S. and Di, Y.-B. (eds) Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 1990/3: p. 23-29
- IFOAM (1995)  
IFOAM conference on food, culture, trade and the environment, 19-22 July 1995, Seoul, Korea.

- Khan, A.M., Saxena, S.K. and Siddiqi, Z.A. (1971) Efficacy of *Tagetes erecta* in reducing root infesting nematodes of tomato and okra. *Indian Phytopathology* 24: p. 166-169.
- Kobayashi, N. (1991) Biological control of soil-borne diseases with VAM fungi and charcoal compost. In: Komada, H, Kiritani, K. and Bay-Peterson, J. (eds) *The biological control of plant diseases. Proceedings of the international seminar Tsukuba, Japan, 17-21 September 1990*: p. 153-160
- Kuwata, H., Saitoh, H. and Shimada, K. (1994) Occurrence of *Fusarium* wilt of tomato, caused by *Fusarium oxysporum* f.sp. *lycopersici* race J2, in Aomori Prefecture, and its control by grafting to rootstocks. *Bulletin of the Aomori agricultural experiment station* 34: p. 1 -10
- Lodha, S. (1995) Soil solarisation, summer irrigation and amendments for the control of *Fusarium oxysporum* f.sp. *cumini* and *Macrophomina phaseolina* in and soils. *Crop protection* 14(3): 215-219
- Lu, MA Chen, Y.H., Lin, M.S. and Chen, H.P. (1992) An experiment using grafting for control of tomato bacterial wilt. *Plant protection* 18(3): p. 25
- Marois, J.J., Johnston, S.A., Dunn, M.T. and Papavizas, G.C. (1982) Biological control of Verticillium wilt of eggplant in the field. *Plant disease* 66: p. 1166-1168
- Morra, L., Mennella, G. and D'Amore, R. (1992) Innesto della melanzana (*Solanum melongena* L.) quale mezzo per la difesa da patogeni tellurici e l'aumento dell resa. 11. Contributo. *Culture Prolette* 21(12): p. 85-93
- Osmelak, J.A. (1990) Status and management of major vegetable pests in Australia. In: Lim, G.S. and Di, Y.-B. (eds) *Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 1990/3*: p. 12-22
- Porter, I.J. and Merriman, P.R. (1985) Evaluation of soil solarisation for control of clubroot of crucifers and white rot of onions in Southeastern Australia. In: Parker, C.A., Rovira, A.D., Moore, K.J. and Wong, P.T.W. (eds) *Ecology and management of soil-borne plant pathogens. American Phytopathological Society St Paul*: p. 282-284
- Prasad, D. and Chawla, M.L. (1991) Importance of environmental factors in management of phyto-nematodes. In: Mohan, 1. (ed) *The fragile environment. New World Environment Series, New Delhi*, 5: p. 115-125
- Prasad, J.S. and Reddy, K.V.S. (1994) Hot water treatment for banana planting material made easier. *Infornusa* 3(2): p. 16
- Purwati, E. and Hanudin (1995) Screening of some tomato genotypes for resistance to bacterial wilt by IBWDN method in Indonesia. *Proceedings of the AVNET 11 midterm workshop, Los Banos, the Philippines*: p. 127-129
- Qiu, W.F. (1990) Part 1: Status and management of major vegetable -pests in the peoples republic of China. In: Lim, G.S. and Di, Y.-B. (eds) *Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 1990/3*: p. 30-35

- Rangaswamy, S.D. and Reddy, P.P. (1993)  
Effect of leaf extracts of trap crops on the growth of tomato and development of *Meloidogyna incognita*. Current nematology 4(1): p. 7-10
- Rao, G.N. (1976)  
Control of nematodes in nursery soils. Plant. Chron. 71: p. 257, 259
- Sariah, M. and Tanaka, K. (1994)  
Effect of flooding on the chilli collar rot pathogen in Malaysian soil. 4th international conference on plant protection in the tropics 28-31 March, Kuala Lumpur, Malaysia: p. 90-91
- Sethi, C.L. and Gaur, H.S. (1986)  
Nematode management: an overview. In: Plant parasitic nematodes of India. Problems and progress. Indian Agric. Res. Inst., New Delhi: p. 424-445
- Takeuchi, S. (1990)  
Status and management of major vegetable pests in Japan. In: Lim, G.S. and Di, Y.-B. (eds) Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 1990/13: p. 46-56
- UNDP (1995)  
UNDP survey on methyl bromide. Draft regional report on South-East Asia and the Pacific region. Draft SEAP report (29 October 1995): pp. 39.
- Vattanatangum, A. (1990)  
Status and management of major vegetable pests in Thailand. In: Lim, G.S. and Di, Y.-B. (eds) Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 19W3: p. 91-95
- Vos, J.G.M. and Sumami, N. (1997)  
Integrated crop management of hot pepper (*Capsicum* spp.) under tropical lowland conditions: effects of mulch on crop performance and production. Journal of Horticultural Science 72(2): In press
- Vos, J.G.M., Nurtika, N. and Sumami, N. (1993)  
An exploratory survey on farmers' practices and management of hot pepper (*Capsicum* spp.) in Java, Indonesia. Journal of Plant Protection in the Tropics 10(2): p. 91-109.
- VU, V.M. (1990)  
Status and management of major vegetable pests in Vietnam. In: Lim, G.S. and Di, Y.-B. (eds) Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 1990/3: p. 96-100